



CENTRE FOR STOCHASTIC GEOMETRY
AND ADVANCED BIOIMAGING

Luis M. Cruz-Orive

**Course on
Recent results on stereology (part II)**



Recent results on stereology

Luis M. Cruz-Orive

*Department of Mathematics, Statistics and Computation
University of Cantabria, E-Santander*

University of DK-Aarhus, 14–15.12.2011.

Part II

1. Computation of the flower area and the wedge volume for an interior pivotal point.

— The pivotal section is convex with smooth boundary of known parametric coordinates.

— The pivotal section is a convex polygon.

2. Uniqueness properties of the invariator. Connections with the nucleator and the surfactor. Computational implications.

3. Open questions, final discussion.

The invariator principle and its applications

Computational problems

- (i). To exploit the flower formula $S(\partial Y) = 4\mathbb{E}A(H_t)$ we have to compute the flower area of any planar convex set.
- (ii). To exploit the wedge formula $V(Y) = 2\pi \mathbb{E}V(W_t)$ we have to compute the mean wedge volume of any planar set.

Both problems (i), (ii) have been solved for an arbitrary planar convex n -gon K in the following two cases:

- (a). Interior pivotal point $O \in K$, (Cruz-Orive LM (2011) *J. Microscopy* 243, 86–102).
- (b). Exterior pivotal point $O \notin K$.

Next we concentrate on case (a).

Flower area of a planar convex set with smooth boundary


Consider a compact convex set $K \subset \mathbb{R}^2$ such that $O \in K^0$, with boundary ∂K of class C^2 admitting the following parametric equations,

$$\partial K = \left\{ (X, Y) \in \mathbb{R}^2 : X = X(t), Y = Y(t), (0 \leq t < 2\pi) \right\}.$$

The parametric equations of the support function h_K , namely of the boundary ∂H_K of the flower H_K of K with respect to O , are the following,

$$\left\{ x(t) = -\frac{Y'(t)}{X'(t)} \cdot y(t), \quad y(t) = \frac{X'(t)Y(t) - X(t)Y'(t)}{X'^2(t) + Y'^2(t)} \cdot X'(t), \quad (0 \leq t < 2\pi) \right\}$$

Hint of Proof. $P(x(t), y(t))$ is the intersection between the tangent to ∂K at the point $M(X(t), Y(t))$ and the normal to this tangent from O . Moreover,



$$A(H_K) = \frac{1}{2} \int_0^{2\pi} h^2(\omega) d\omega = \frac{1}{2} \int_0^{2\pi} h^2(t) \omega'(t) dt = \frac{1}{2} \int_0^{2\pi} \frac{[X'(t)Y(t) - X(t)Y'(t)]^2 [X'(t)Y''(t) - X''(t)Y'(t)]}{[X'^2(t) + Y'^2(t)]^2} dt.$$



Case of an ellipse

Let ∂K be an ellipse of principal semiaxes $0 < m \leq M < \infty$, and let $(x_0, y_0) \in K$ represent the rectangular coordinates of the pivotal point with respect to the ellipse centre. Shift the origin to this pivotal point. Then,

$$\partial K : \begin{cases} X(t) = M \cos t - x_0, \\ Y(t) = m \sin t - y_0, \end{cases} \quad (0 \leq t < 2\pi).$$

$$A(H_K) = \frac{\pi}{2} (M^2 + m^2 + r^2),$$

where $r^2 := x_0^2 + y_0^2$.




Flowers of identical areas if r is fixed

Flower area of a convex polygon

A pivotal section of a convex particle is a.s. convex. It can be approximated by a convex n -gon with the required accuracy for sufficiently large n .


Consider a pivotal polygonal section of a convex polyhedron.



A useful representation of the flower of a convex polygon K :

$$H_K = \bigcup_{z \in \partial K} B_2(z/2, \|z\|/2), \quad (O \in K^0).$$

P Calka's formula for the flower area of a convex polygon



$$A(H_K) = \frac{1}{8} \sum_{k=1}^n R_{k+1}^2 \psi(Z_k, Z_{k+1}, F_{k+1}),$$

$\{(r_i, \varphi_i)\} :=$ polar coords. of the vertices, $\{(p_i, \theta_i)\} :=$ normal coords. of the sides,

$R := (\{r_i\}_1^n, \{r_i\}_1^n)$, $F := (\{\varphi_i\}_1^n, \{\varphi_i\}_1^n)$, $Z_i := (\{\theta_i\}_1^n, \{\theta_i + 2\pi\}_1^n)$,

$\psi(\omega_1, \omega_2, \varphi) := \sin(2(\omega_2 - \varphi)) - \sin(2(\omega_1 - \varphi)) + 2(\omega_2 - \omega_1)$,

$v_1(x_1, y_1)$ is the rightmost vertex.

(Calka (2003) AAP 35, 27–46, Calka (2009) personal communication).

P Calka's formula: Hint of proof

When K is a convex polygon its flower H_K is

$$H_K = \bigcup_{k=1}^n B_2((r_k/2, \varphi_k), r_k/2),$$

namely the union of the n disks whose diameters are the radius vectors of the vertices. Consequently,

$$h_K(\omega) = r_{k+1} \cos(\omega - \varphi_{k+1}), \quad \omega \in [\theta_k, \theta_{k+1}),$$

and,

$$\begin{aligned} A(H_K) &= \frac{1}{2} \sum_{k=1}^n \int_{Z_k}^{Z_{k+1}} h_K^2(\omega) d\omega \\ &= \frac{1}{2} \sum_{k=1}^n \int_{Z_k}^{Z_{k+1}} R_{k+1}^2 \cos^2(\omega - \varphi_{k+1}) d\omega, \end{aligned}$$


from which the result follows

Mean wedge volume of a convex polygon (i)

$$V(Y) = 2\pi \mathbb{E}\{V(W_t)\}, \quad V(W_t) := \mathbb{E}V\{W(\omega; t)|t\}$$

Purpose: To find an exact formula for $V(W_t)$.

First step: For fixed $t \in \mathbb{S}_+^2$ and for a given $\omega \in [0, 2\pi)$, compute $V\{W(\omega; t)\}$ by a simplicial decomposition:




$$W(\omega) := W(\omega; t)$$

Mean wedge volume of a convex polygon (ii)


Second step: Compute

$$\mathbb{E}V\{W(\omega)\} = \frac{1}{2\pi} \int_0^{2\pi} V\{W(\omega)\} d\omega.$$


The main task is to identify the integration subranges. This is equivalent to identifying the vertices of the subpolygon $K_+(\omega)$ for each $\omega \in [0, 2\pi)$.



Convex n -gon




Subpolygon identification problem



Subrange identification problem

Mean wedge volume of a convex polygon (iii)

A tool to solve the identification problems: The 'extended polygon'.



'Extended polygon' K_E of K
($2n$ 'vertices')

No identification problems anymore

Mean wedge volume of a convex polygon (iv): Exact formula

The mean wedge volume associated with a convex n -gon K whose interior contains the origin can be expressed as follows,

$$\begin{aligned}
 2\pi \mathbb{E}_\omega V\{W(\omega)\} &= \frac{1}{6} \sum_{k=1}^{2n} \left\{ P_k R_{k+1}^2 \psi_1(F_k, F_{k+1}, F_{k+1}, Z_k) \right. \\
 &\quad + \sum_{i=k+1}^{k+n-1} P_i L_i \cdot [R_i \psi_2(F_k, F_{k+1}, F_i) + R_{i+1} \psi_2(F_k, F_{k+1}, F_{i+1})] \\
 &\quad \left. - P_{k+n} R_{k+n}^2 \psi_1(F_k, F_{k+1}, F_{k+n}, Z_{k+n}) \right\},
 \end{aligned}$$

where the angle vectors F, Z and the length vectors R, P, L are $4n$ -vectors defined on the extended polygon, whereas

$$\begin{aligned}
 \psi_1(\omega_1, \omega_2, \phi, \theta) &:= \sin(2\phi - \theta - \omega_2) - \sin(2\phi - \theta - \omega_1) \\
 &\quad + \cos^2(\phi - \theta) \cdot \log \left(\frac{\tan((\omega_2 - \theta)/2 + \pi/4)}{\tan((\omega_1 - \theta)/2 + \pi/4)} \right), \\
 \psi_2(\omega_1, \omega_2, \phi) &:= \cos(\omega_2 - \phi) - \cos(\omega_1 - \phi). \quad \square
 \end{aligned}$$

Mean wedge volume of a convex polygon (iv): Approximate formula for large n

Monte Carlo integration with N steps:

$$2\pi \hat{\mathbb{E}}_{\omega} V\{W(\omega)\} = \frac{2\pi}{N} \sum_{i=1}^N V\{W(\omega_i)\}, \quad \omega_i = (U + i - 1) \cdot \frac{2\pi}{N} + \frac{\pi}{2}, \quad i = 1, 2, \dots, N, \quad U \sim \text{UR}(0, 1).$$

An unbiased estimator of the mean wedge volume associated with a convex n -gon K whose interior contains the origin can be computed as follows,

$$\begin{aligned} 2\pi \hat{\mathbb{E}}_{\omega} V\{W(\omega)\} &= \frac{2\pi}{N} \cdot \frac{1}{6} \sum_{i=1}^N \left\{ P_{k(i)} R_{k(i)+1}^2 \psi\left(\omega_i, F_{k(i)+1}, Z_{k(i)}\right) \right. \\ &\quad + \sum_{j=k(i)+1}^{k(i)+n-1} P_j L_j \cdot [R_j \cos(\omega_i - F_j) + R_{j+1} \cos(\omega_i - F_{j+1})] \\ &\quad \left. - P_{k(i)+n} R_{k(i)+n}^2 \psi\left(\omega_i, F_{k(i)+n}, Z_{k(i)+n}\right) \right\}, \end{aligned}$$


where the angle vectors F, Z and the length vectors R, P, L are defined as before; the index $k(i)$ is such that the axis $L_{1[0]}(\omega_i)$ hits the sides $S_{k(i)}$ and $S_{k(i)+n}$ of the extended polygon, and

$$\psi(\omega, \phi, \theta) := \frac{\cos^2(\omega - \phi)}{\sin(\omega - \theta)}.$$


Furthermore,

$$\text{Var}\left\{\hat{\mathbb{E}}_{\omega} V\{W(\omega)\}\right\} = O(N^{-\alpha}), \quad \alpha \in [2, 4].$$


The mean wedge volume of a disk



Ball and pivotal section



Pivotal section



Wedge

An UE of the volume of the ball is,



$$\widehat{V}(Y; \lambda) = 2\pi E_{\omega} V\{W(\omega)\} = \frac{8}{9}r^3 \left[(7 + \lambda^2)E(\lambda^2) - 4(1 - \lambda^2)K(\lambda^2) \right], \quad \lambda = l/r \in (0, 1),$$

where K , E represent the complete elliptic integrals of the first and second kinds, respectively. Note that $\widehat{V}(Y; 0) = (4/3)\pi r^3$.

Approximating the disk by a regular polygon of $n = 5000$ vertices, the Monte Carlo approximation yielded 6 exact digits of $\widehat{V}(Y; \lambda)$ for $\lambda = 0.1, 0.3, 0.7, 0.9$, with $N = 20$ steps. Real computing time: 2s. With the exact formula: 102s.

Mean volume weighted surface area and volume of cemented carbide grains (i)

Material. Cemented carbides constitute a group of powder metallurgical materials produced by sintering, and characterized by a very high hardness and wear resistance. Their microstructure consists of hard tungsten carbide (WC) grains embedded in a ductile cobalt (Co) matrix.



$$\mathbb{E}_W(X) = \frac{\mathbb{E}_N(WX)}{\mathbb{E}_N(W)}, \text{ here we want } \mathbb{E}_V(S), \mathbb{E}_V(V).$$

Mean volume weighted surface area and volume of cemented carbide grains (ii)

Let $1 \leq p < \infty$ be the number of test points hitting grains, and let s_i, v_i denote unbiased estimators of the surface area and the volume of the grain hit by the i -th test point, respectively, ($i = 1, 2, \dots, p$). Because the mentioned grain is volume-weighted, the estimators:

$$\bar{s}_V = \frac{1}{p} \sum_{i=1}^p s_i = \frac{4}{p} \sum_{i=1}^p F_i, \quad \text{where } F_i := \text{flower area},$$

$$\bar{v}_V = \frac{1}{p} \sum_{i=1}^p v_i = \frac{2\pi}{p} \sum_{i=1}^p W_i, \quad \text{where } W_i := \text{mean wedge volume},$$

are ratio unbiased for the target volume weighted means $\mathbb{E}_V S$ and $\mathbb{E}_V V$, respectively. In shorthand,

$$\bar{s}_V = 4\bar{F},$$

$$\bar{v}_V = 2\pi\bar{W}.$$

Mean volume weighted surface area and volume of cemented carbide grains (iii)

Results for the cemented carbide grains.

The 1997 results were reported in Karlsson & C-O (1997). For the “surfator” and “PSI” (point sampled intercepts) methods, see the mentioned study.

Quadrat	# polygons	$\bar{A}, \mu\text{m}^2$	$\bar{s}_V = 4\bar{F}, \mu\text{m}^2$	$\bar{v}_V = 2\pi\bar{W}, \mu\text{m}^3$
1	20	21.24	144.9	135.5
2	17	32.36	209.3	233.6
3	16	24.76	160.1	160.3
4	17	21.44	142.5	124.9
Pool	70	24.80 (10.4%)	163.4 (9.5%)	162.4 (15.0%)
Results of 1997			$\bar{s}_V \mu\text{m}^2$, surfator	$\bar{v}_V \mu\text{m}^3$, PSI
			154.5 (4.4%)	161.3(23.7%)

Uniqueness properties of the invariator

The mean wedge volume is the averaged nucleator

Bounded particle: $Y \subset B_3 \subset \mathbb{R}^3$.

Mean wedge volume representation of the particle volume:

$$V(Y) = 2\pi \mathbb{E}_t \mathbb{E}_\omega V\{W(\omega; t)\}.$$

Nucleator representation (to simplify the notation suppose for the moment that Y is star shaped with respect to an interior pivotal point O):

$$V(Y) = \frac{4\pi}{3} \mathbb{E}_t \mathbb{E}_\varphi \left\{ \rho_+^3(\varphi; t) \right\}.$$

Proposition 1. *For any pivotal section of a particle with respect to an interior pivotal point, namely for each orientation $t \in \mathbb{S}_+^2$ of the pivotal plane, the following identity holds,*


$$\mathbb{E}_\omega V\{W(\omega; t)\} = \frac{2}{3} \mathbb{E}_\varphi \left\{ \rho_+^3(\varphi; t) \right\},$$

that is, the mean wedge volume coincides with 2/3 times the averaged third power of the nucleator ray length.

Proof. For each $t \in \mathbb{S}_+^2$ and each $\omega \in [0, 2\pi)$,

$$V\{W(\omega; t)\} = \int_0^{h(\omega; t)} l(r, \omega; t) r dr.$$


Consider a point of unsigned abscissa u along the p -line with respect to an arbitrary origin on this line. Then clearly,



$$l(r, \omega; t) = \int_{(Y \cap L_2^3(0, t)) \cap L_1^2(r, \omega; t)} du,$$

and the length elements du and dr are orthogonal. Thus $dx := du dr$ is the area element at a point x in the pivotal plane. Let (ρ, φ) , $(\rho > 0, 0 \leq \varphi < 2\pi)$ denote the polar coordinates of x with respect to the fixed axis Ox_1 .

Then $r = \rho |\cos(\omega - \varphi)|$ and we can write,



$$\begin{aligned}
 \mathbb{E}_\omega V\{W(\omega; t)\} &= \frac{1}{2\pi} \int_0^\pi d\omega \int_{Y \cap L_2^3(0,t)} \rho |\cos(\omega - \varphi)| dx \\
 &= \frac{1}{2\pi} \int_{Y \cap L_2^3(0,t)} \rho dx \int_0^\pi |\cos(\omega - \varphi)| d\omega \\
 &= \frac{1}{\pi} \int_{Y \cap L_2^3(0,t)} \rho dx \\
 &= \frac{1}{\pi} \int_0^{2\pi} d\varphi \int_0^{\rho_+(\varphi; t)} \rho^2 d\rho \\
 &= \frac{2}{3} \mathbb{E}_\varphi \left\{ \rho_+^3(\varphi; t) \right\}.
 \end{aligned}$$

Computation of the mean wedge volume defined on a convex polygon via the averaged nucleator: a vast improvement on Cruz-Orive (2011)

Corollary 1. *For a convex n -gon $K \subset \mathbb{R}^2$ the mean volume of the wedge set $W_K(\omega)$ defined on K with respect to an arbitrary pivotal point in its plane may be expressed as follows,*

$$2\pi \mathbb{E}_\omega V\{W_K(\omega)\} = \frac{1}{3} \sum_{k=1}^n p_k^3 \cdot \psi(\varphi_k, \varphi_{k+1}, \theta_k),$$

where

$$\begin{aligned} \psi(\omega_1, \omega_2, \theta) &:= \frac{\sin(\omega_2 - \theta)}{\cos^2(\omega_2 - \theta)} - \frac{\sin(\omega_1 - \theta)}{\cos^2(\omega_1 - \theta)} \\ &\quad + \log \left(\frac{\tan((\omega_2 - \theta)/2 + \pi/4)}{\tan((\omega_1 - \theta)/2 + \pi/4)} \right). \end{aligned}$$

If $O \notin K^\circ$, then K must be contained in the upper half plane.

Proof. For $k = 1, 2, \dots, n$ let $\rho_k(\omega)$, $\omega \in [\varphi_k, \varphi_{k+1})$ denote the radial function of ∂K . Bearing in mind that $\rho_k(\omega) \cos(\omega - \theta_k) = p_k$, by virtue of Proposition 1 we have,

$$2\pi \mathbb{E}_\omega V\{W(\omega; t)\} = \frac{2}{3} \sum_{k=1}^n \int_{\varphi_k}^{\varphi_{k+1}} \rho_k^3(\omega) d\omega = \frac{2}{3} \sum_{k=1}^n p_k^3 \int_{\varphi_k}^{\varphi_{k+1}} \frac{d\omega}{\cos^3(\omega - \theta_k)}.$$

On the other hand,


$$\int \frac{dx}{\cos^3 x} = \frac{1}{2} \frac{\sin x}{\cos^2 x} + \frac{1}{2} \log \tan \left(\frac{x}{2} + \frac{\pi}{4} \right) + C,$$

whereby Corollary 1 follows.

Checking the new mean wedge volume formula


Assume that the particle $Y \subset \mathbb{R}^3$ is a finite ball. The following exact results have been checked numerically with the formula.

Case of an interior pivotal point



$$2\pi E_{\omega} V\{W(\omega)\} = \frac{8}{9} r^3 \left[(7 + \lambda^2) E(\lambda^2) - 4(1 - \lambda^2) K(\lambda^2) \right], \quad \lambda := l/r \in (0, 1)$$

Case of an exterior pivotal point



$$2\pi E_{\omega} V\{W(\omega)\} = \frac{8}{9\lambda} \left[\lambda^2 (7 + \lambda^2) E(\lambda^{-2}) - (\lambda^2 + 3)(\lambda^2 - 1) K(\lambda^{-2}) \right], \quad \lambda := l/r > 1$$


The integrated surfactor is the flower area

For a convex particle $Y \subset \mathbb{R}^3$ containing the pivotal point, the flower formula for the particle surface area reads,

$$S(\partial Y) = 4 \mathbb{E}_t \{ A(H_t) \} = 2 \mathbb{E}_t \left\{ \int_0^{2\pi} h^2 d\omega \right\}$$

Alternatively, the surfactor representation is,

$$S(\partial Y) = 4\pi \mathbb{E}_t \mathbb{E}_\varphi \left\{ \rho^2 (1 + \alpha \tan \alpha) \right\} = 2 \mathbb{E}_t \left\{ \int_0^{2\pi} \rho^2 (1 + \alpha \tan \alpha) d\varphi \right\}$$



The next proposition states that the two expressions in curly brackets coincide for each $t \in \mathbb{S}^2$.

Case of a convex particle with smooth boundary

Proposition 2. For each pivotal section of a convex particle with boundary of class C^2 containing the pivotal point, the following identity holds,


$$\int_0^{2\pi} \rho^2(1 + \alpha \tan \alpha) d\varphi = \int_0^{2\pi} h^2 d\omega.$$

Lemma 1. The flower area $A(H_K)$ of a planar convex set K with boundary of class C^2 containing the origin may be expressed as follows,

$$A(H_K) = A(K) + \frac{1}{2} \int_0^{2\pi} \rho^2 \sin^2 \alpha d\omega,$$


where the angle α is defined as in Eq. 24. Equivalently,

$$\int_0^{2\pi} h^2 d\omega = \int_0^{2\pi} \rho^2 d\varphi + \int_0^{2\pi} \rho^2 \sin^2 \alpha d\omega.$$



Proof 1 of Lemma 1. Clear from the figure.

Proof 2 of Lemma 1. The ordinary area element dA in polar coordinates is $(1/2)\rho^2 d\varphi$. However, for the present purposes we need a different representation. The area element may be regarded as an elementary triangle OMM' of height $\rho \cos \alpha$ and base $\sin \alpha d\rho + \rho \cos \alpha d\varphi$, see the figure.



Therefore,

$$dA = (1/2)\rho \sin \alpha \cos \alpha d\rho + (1/2)\rho^2 \cos^2 \alpha d\varphi = (1/2) \rho \rho'_\alpha \sin \alpha \cos \alpha d\alpha + (1/2) \rho^2 \cos^2 \alpha d\varphi,$$

where $\rho'_\alpha := d\rho/d\alpha$. Thus,

$$2A(K) = \int_0^{2\pi} \rho \rho'_\alpha \sin \alpha \cos \alpha d\alpha + \int_0^{2\pi} \rho^2 \cos^2 \alpha d\varphi.$$


Integration by parts with $u = \sin \alpha \cos \alpha$ and $dv = \rho \rho'_\alpha d\alpha$ yields,

$$\int_0^{2\pi} \rho \rho'_\alpha \sin \alpha \cos \alpha d\alpha = -\frac{1}{2} \int_0^{2\pi} \rho^2 (\cos^2 \alpha - \sin^2 \alpha) d\alpha.$$

Recalling that $\alpha = \varphi - \omega$ and $\rho \cos \alpha = h$, simplification leads to the required result.

Proof of Proposition 2. From the right hand side figure below we obtain the key relation

$$\tan \alpha = \frac{d\rho}{\rho d\varphi} = \frac{\rho'_\varphi}{\rho}.$$



Therefore,

$$\int_0^{2\pi} \rho^2 \alpha \tan \alpha d\varphi = \int_0^{2\pi} \rho \rho'_\varphi \alpha d\varphi.$$

Integration by parts with $u = \alpha$ and $dv = \rho \rho'_\varphi d\varphi$ yields,

$$\int_0^{2\pi} \rho^2 \alpha \tan \alpha d\varphi = -\frac{1}{2} \int_0^{2\pi} \rho^2 d\alpha = -\frac{1}{2} \int_0^{2\pi} \rho^2 d\varphi + \frac{1}{2} \int_0^{2\pi} \rho^2 (\sin^2 \alpha + \cos^2 \alpha) d\omega.$$

Bearing in mind that $\rho \cos \alpha = h$

$$\begin{aligned}
 \int_0^{2\pi} \rho^2(1 + \alpha \tan \alpha) d\varphi &= \int_0^{2\pi} \rho^2 d\varphi - \frac{1}{2} \int_0^{2\pi} \rho^2 d\varphi + \frac{1}{2} \int_0^{2\pi} \rho^2 \sin^2 \alpha d\omega + \frac{1}{2} \int_0^{2\pi} h^2 d\omega \\
 &= \frac{1}{2} \int_0^{2\pi} h^2 d\omega - \frac{1}{2} \int_0^{2\pi} \rho^2 \sin^2 \alpha d\omega + \frac{1}{2} \int_0^{2\pi} \rho^2 \sin^2 \alpha d\omega + \frac{1}{2} \int_0^{2\pi} h^2 d\omega \quad (\text{Lemma 1}) \\
 &= \int_0^{2\pi} h^2 d\omega. \quad \square
 \end{aligned}$$

CASE OF A CONVEX POLYHEDRAL PARTICLE

Proposition 2 has been proved under the assumption that the particle Y has a unique tangent plane at every point of its boundary. Because this property may look restrictive, here we assume that Y is a convex polyhedron. Thus, every pivotal section is almost surely a convex polygon.

Proposition 3. *Proposition 2 holds also for any convex n -gon with an interior pivotal point.*

Proof. Recalling that $\rho_k(\omega) \cos(\omega - \theta_k) = p_k$, we can write,

$$\int_0^{2\pi} \rho^2(1 + \alpha \tan \alpha) d\varphi = \sum_{k=1}^n \int_{\varphi_k}^{\varphi_{k+1}} \frac{p_k^2}{\cos^2 \alpha} (1 + \alpha \tan \alpha) d\varphi,$$

where $\alpha := \varphi - \theta_k$ in the preceding integrand. On the other hand, by Calka's formula,

$$\int_0^{2\pi} h^2 d\omega = \frac{1}{4} \sum_{k=1}^n r_{k+1}^2 [\zeta(\theta_{k+1} - \varphi_{k+1}) - \zeta(\theta_k - \varphi_{k+1})],$$

$$\zeta(x) := 2x + \sin(2x).$$


Thus, the right hand sides of the preceding two equations must coincide. This is readily verified on substituting the following results

$$\int \frac{1 + x \tan x}{\cos^2 x} dx = \frac{1}{2} \tan x + \frac{1}{2} \frac{x}{\cos^2 x} + C,$$


$$p_k = r_k \cos(\theta_k - \varphi_k) = r_{k+1} \cos(\theta_k - \varphi_{k+1}),$$

into the right hand side of the former equation, and simplifying.


Flower of a convex set with exterior pivotal point: A glimpse



Support function $h(\omega)$ with exterior pivotal point O



Support set (flower) H_K




Unwrapped graph of $h(\omega)$

$$h(\omega) = x_0 \cos \omega + y_0 \sin \omega + h_0(\omega), \quad \omega \in [0, 2\pi),$$

$$A(H_K) = \frac{1}{2} \int_0^{2\pi} h(\omega) |h(\omega)| d\omega.$$

Flowers of an ellipse with exterior pivotal point







$l = 3, M = 1, m = 1/\sqrt{2}$

$A(H_K) = 10.9869$

10.7190

10.2063

9.9597



With an exterior pivotal point, the flower area of an ellipse depends not only on M, m, l , but also on α .

References

- Calka P (2003) *Adv. Appl. Prob.* **35**, 863–870.
- Calka P (2009) Personal communication.
- C-O (2005) *J Microsc* **219**, 18–28.
- C-O (2008) *Image Anal Stereol* **27**, 17–22.
- C-O (2009) *Image Anal Stereol* **28**, 63–67.
- C-O (2011) *J Microsc* **243**, 86–102.
- C-O (2011) *Int Report 1/2011*, MATESCO, UC.
- C-O, Ramos-Herrera ML & Artacho-Pérula (2010) *J Microsc* **240**, 94–110.
- Gual-Arnau X & C-O (2009) *Diff Geom Appl* **27**, 124–128.
- Gual-Arnau X., C-O & Nuño-Ballesteros JJ (2010) *Adv Appl Math* **44**, 298–308.
- Gundersen HJG (1988) *J Microsc* **151**, 3–21.
- Jensen EBV (1991) *Ann Inst Statist Math* **43**, 455–468.
- Jensen EB and Gundersen HJG (1989) *J Microsc* **153**, 249–267.
- Jensen EBV and Rataj J (2008) *Adv Appl Math* **41**, 530–560.
- Karlsson, LM & C-O (1997) *J Microsc* **186**, 121–132.
- Varga O (1935) *Math Zeitschrift* **40**, 387–405.